

Table: Statistics for variable (area) and region (cna-n). Model N is the average 34-member objects.

Statistic	CRU	PRISM	CESMLE(C)	CESMLE(S1)	CESMLE(S2)	CESMLE(S3)
N	17	18	15.15	15.35	13.47	10.15
Mean	14	13	11.80	6.70	4.97	4.12
SD	34	31	31.19	15.94	8.60	7.28
10 th P	1.00	1.00	1.00	1.00	1.00	1.10
Q1	1.00	1.22	1.70	1.60	1.60	1.60
Median	1.7	1.8	2.20	2.50	2.35	2.30
Q2	3.20	3.28	4.50	4.80	4.80	3.80
90 th P	37.62	38.48	18.36	9.90	9.26	7.12
¹ Normality p	< 0.01 *	< 0.01 *	< 0.01 *	< 0.01 *	< 0.01 *	< 0.01 *
² Welch p		0.98	0.82 0.83	< 0.01 *	< 0.01 *	< 0.01 *
³ Tukey p		1.00	1.00 1.00	< 0.01 *	< 0.01 *	< 0.01 *
⁴ WMW p		0.53	0.04 0.14	0.75	0.84	0.23

All tests significant at $\alpha = 0.05$.

¹ Shapiro-Wilk Normality Test. * denotes the distribution is not a normal distribution.

² Welch Two-Sample T-Test. * denotes the paired sample distribution means are significantly different.

³ Tukey Honestly Significant Difference Test. * denotes the paired sample distribution means are significantly different.

⁴ Wilcoxon Mann-Whitney U Test. * denotes the sample distributions are significantly different.