

Table: Statistics for variable (axis-ang) and region (cna-n). Model N is the average 34-member objects.

Statistic	CRU	PRISM	CESMLE(C)	CESMLE(S1)	CESMLE(S2)	CESMLE(S3)
N	19	23	15.41	9.18	5.41	4.35
Mean	0.69	8.6	5.80	2.78	2.72	2.86
SD	35	30	28.34	26.53	24.23	29.30
10 th P	-33.55	-6.22	-19.45	-26.40	-10.36	-12.50
Q1	-1.69	0.00	0.00	0.00	-4.06	-7.00
Median	0	0	0.00	0.00	0.00	0.00
Q2	4.66	2.66	12.95	4.93	0.00	0.00
90 th P	37.58	41.00	42.61	35.89	31.60	45.00
¹ Normality p	0.01*	<0.01*	<0.01*	<0.01*	<0.01*	<0.01*
² Welch p		0.44	0.53 0.66	0.12	0.16	0.28
³ Tukey p		0.94	0.97 1.00	0.64	0.78	0.86
⁴ WMW p		0.65	0.63 0.93	0.11	0.02*	<0.01*

All tests significant at $\alpha = 0.05$.

¹ Shapiro-Wilk Normality Test. * denotes the distribution is not a normal distribution.

² Welch Two-Sample T-Test. * denotes the paired sample distribution means are significantly different.

³ Tukey Honestly Significant Difference Test. * denotes the paired sample distribution means are significantly different.

⁴ Wilcoxon Mann-Whitney U Test. * denotes the sample distributions are significantly different.