

Table: Statistics for variable (axis-ang) and region (cna-s). Model N is the average 34-member objects.

Statistic	CRU	PRISM	CESMLE(C)	CESMLE(S1)	CESMLE(S2)	CESMLE(S3)
N	26	17	12.62	11.12	10.59	8.59
Mean	8.1	-3.3	17.09	16.06	14.17	15.45
SD	56	30	43.46	44.24	44.19	49.40
10 th P	-60.08	-50.22	-42.66	-45.96	-46.97	-64.32
Q1	-30.31	0.00	-1.13	0.00	-6.61	-0.56
Median	0	0	10.94	7.69	2.16	9.32
Q2	59.19	6.79	49.08	46.25	45.00	55.74
90 th P	90.00	27.89	77.71	82.75	76.00	85.72
¹ Normality p	0.04*	0.03*	<0.01*	<0.01*	<0.01*	<0.01*
² Welch p	0.39		0.43 0.02*	0.74	0.35	0.65
³ Tukey p	0.97		0.92 0.45	1.00	0.95	1.00
⁴ WMW p	0.91		0.28 0.05	0.79	0.30	0.97

All tests significant at $\alpha = 0.05$.

¹ Shapiro-Wilk Normality Test. * denotes the distribution is not a normal distribution.

² Welch Two-Sample T-Test. * denotes the paired sample distribution means are significantly different.

³ Tukey Honestly Significant Difference Test. * denotes the paired sample distribution means are significantly different.

⁴ Wilcoxon Mann-Whitney U Test. * denotes the sample distributions are significantly different.