

Table: Statistics for variable (intensity-25) and region (cna-s). Model N is the average 34-member objects.

Statistic	CRU	PRISM	CESMLE(C)	CESMLE(S1)	CESMLE(S2)	CESMLE(S3)
N	26	17	12.62	11.12	10.59	8.59
Mean	-1.4	-1.3	-1.34	-1.38	-1.37	-1.39
SD	0.2	0.18	0.20	0.24	0.25	0.24
10 th P	-1.61	-1.54	-1.65	-1.74	-1.75	-1.69
Q1	-1.42	-1.37	-1.42	-1.47	-1.47	-1.45
Median	-1.3	-1.3	-1.28	-1.32	-1.30	-1.34
Q2	-1.25	-1.19	-1.19	-1.21	-1.20	-1.24
90 th P	-1.15	-1.11	-1.14	-1.14	-1.13	-1.17
¹ Normality p	< 0.01 *	0.18	< 0.01 *	< 0.01 *	< 0.01 *	< 0.01 *
² Welch p		0.35	0.52 0.53	0.01 *	0.03 *	< 0.01 *
³ Tukey p		0.98	0.99 1.00	0.11	0.26	0.02 *
⁴ WMW p		0.33	0.27 0.71	0.01 *	0.10	< 0.01 *

All tests significant at $\alpha = 0.05$.

¹ Shapiro-Wilk Normality Test. * denotes the distribution is not a normal distribution.

² Welch Two-Sample T-Test. * denotes the paired sample distribution means are significantly different.

³ Tukey Honestly Significant Difference Test. * denotes the paired sample distribution means are significantly different.

⁴ Wilcoxon Mann-Whitney U Test. * denotes the sample distributions are significantly different.