

Table: Statistics for variable (axis-ang) and region (cna-s). Model N is the average 34-member objects.

Statistic	CRU	PRISM	CESMLE(C)	CESMLE(S1)	CESMLE(S2)	CESMLE(S3)
N	20	24	11.24	8.26	4.94	4.12
Mean	6.6	2.9	14.25	16.60	13.61	9.21
SD	45	40	40.31	41.40	44.15	47.62
10 th P	-47.89	-46.67	-41.05	-36.32	-45.00	-62.18
Q1	-2.80	0.00	0.00	0.00	0.00	0.00
Median	0	0	4.84	0.00	0.00	0.00
Q2	29.05	12.54	44.82	45.00	45.00	45.00
90 th P	83.30	57.04	69.67	82.53	90.00	79.69
¹ Normality p	0.03*	0.01*	<0.01*	<0.01*	<0.01*	<0.01*
² Welch p	0.78		0.46 0.19	0.46	0.87	0.27
³ Tukey p	1.00		0.97 0.80	0.98	1.00	0.84
⁴ WMW p	0.99		0.28 0.17	0.67	0.46	0.24

All tests significant at $\alpha = 0.05$.

¹ Shapiro-Wilk Normality Test. * denotes the distribution is not a normal distribution.

² Welch Two-Sample T-Test. * denotes the paired sample distribution means are significantly different.

³ Tukey Honestly Significant Difference Test. * denotes the paired sample distribution means are significantly different.

⁴ Wilcoxon Mann-Whitney U Test. * denotes the sample distributions are significantly different.