Table: Statistics for variable (axis-ang) and region (ena-s). Model N is the average 34-member objects.

Statistic	CRU	PRISM	CESMLE(C)	CESMLE(S1)	CESMLE(S2)	CESMLE(S3)
N	12	9	10.50	10.18	8.56	8.29
Mean	13	22	18.60	11.87	1.30	-11.75
SD	48	35	46.15	41.89	50.96	50.72
10^{th}P	-42.44	0.00	-45.00	-37.69	-79.45	-80.49
Q1	-4.84	0.00	-0.00	0.00	-19.96	-65.15
Median	0	0	13.06	0.00	0.00	0.00
Q2	36.04	34.22	48.30	40.94	32.16	0.00
90^{th}P	86.56	78.21	90.00	76.47	90.00	79.77
[1]Normality p	0.50	<**0.01***	<**0.01***	<**0.01***	<**0.01***	<**0.01***
[2]Welch p	0.65		0.71 0.80	**0.04***	<**0.01***	<**0.01***
[3]Tukey p	1.00		1.00 1.00	0.41	<**0.01***	<**0.01***
[4]WMW p	0.71		0.60 0.88	**0.02***	<**0.01***	<**0.01***

All tests significant at $\alpha = 0.05$.
[1] Shapiro-Wilk Normality Test. * denotes the distribution is not a normal distribution.
[2] Welch Two-Sample T-Test. * denotes the paired sample distribution means are significantly different.
[3] Tukey Honestly Significant Difference Test. * denotes the paired sample distribution means are significantly different.
[4] Wilcoxon Mann-Whitney U Test. * denotes the sample distributions are significantly different.