

Table: Statistics for variable (axis-ang) and region (ena-s). Model N is the average 34-member objects.

Statistic	CRU	PRISM	CESMLE(C)	CESMLE(S1)	CESMLE(S2)	CESMLE(S3)
N	16	13	9.00	5.09	1.74	0.71
Mean	11	2.9	14.59	16.38	8.83	22.69
SD	49	43	46.68	43.01	51.10	49.38
10 th P	-43.88	-51.21	-52.06	-43.16	-59.02	-29.65
Q1	-24.76	0.00	0.00	0.00	-23.68	0.00
Median	0	0	0.00	0.00	0.00	0.00
Q2	45.61	0.00	45.00	45.00	45.00	79.52
90 th P	88.13	58.20	90.00	83.46	90.00	90.00
¹ Normality p	0.14	0.06	< 0.01 *	< 0.01 *	< 0.01 *	< 0.01 *
² Welch p		0.62	0.80 0.35	0.67	0.42	0.44
³ Tukey p		1.00	1.00 0.95	1.00	0.95	0.96
⁴ WMW p		0.80	0.57 0.26	0.85	0.32	0.67

All tests significant at $\alpha = 0.05$.

¹ Shapiro-Wilk Normality Test. * denotes the distribution is not a normal distribution.

² Welch Two-Sample T-Test. * denotes the paired sample distribution means are significantly different.

³ Tukey Honestly Significant Difference Test. * denotes the paired sample distribution means are significantly different.

⁴ Wilcoxon Mann-Whitney U Test. * denotes the sample distributions are significantly different.