

Table: Observation and model statistics for area, organized by region. Model N is the average 34-member objects.

Obs/Model	Statistic	wna_n	wna_s	cna_n	cna_s	ena_n	ena_s
Cru	N	—	—	—	—	—	—
	Mean	—	—	—	—	—	—
	SD	—	—	—	—	—	—
	Median	—	—	—	—	—	—
Prism	N	—	—	—	—	—	—
	Mean	—	—	—	—	—	—
	SD	—	—	—	—	—	—
	Median	—	—	—	—	—	—
CESMLE(C)	N	0.09	0.03	0.03	—	—	0.06
	Mean	1.63	—	—	—	—	2.05
	SD	0.55	—	—	—	—	—
	Median	1.90	—	—	—	—	—
CESMLE(S1)	N	0.09	0.15	0.24	0.18	—	—
	Mean	2.20	2.66	1.39	2.63	—	—
	SD	1.51	0.89	0.51	1.62	—	—
	Median	1.70	2.50	1.20	2.15	—	—
CESMLE(S2)	N	0.09	0.12	0.06	0.18	—	—
	Mean	2.00	1.35	2.40	1.50	—	—
	SD	0.10	0.33	—	0.61	—	—
	Median	2.00	1.30	—	1.35	—	—
CESMLE(S3)	N	0.12	0.15	0.12	0.24	—	0.06
	Mean	3.52	6.88	1.45	1.52	—	1.00
	SD	2.50	5.45	0.53	0.64	—	—
	Median	2.75	5.20	1.40	1.40	—	—
Cru-Prism		—	—	—	—	—	—
C-Cru		—	—	—	—	—	—
C-Prism	¹ WMW-p	—	—	—	—	—	—
S1-C		1.00	—	—	—	—	—
S2-C		0.26	—	—	—	—	—
S3-C		0.48	—	—	—	—	—

¹ Wilcoxon Mann-Whitney U Test. Sample distributions are significantly different (*) at $\alpha = 0.05$.

² CESMLE time periods: (C) 1981-2010, (S1) 2011-2040, (S2) 2041-2070, and (S3) 2071-2099.