

Table: Statistics for variable (axis-ang) and region (wna-n). Model N is the average 34-member objects.

Statistic	CRU	PRISM	CESMLE(C)	CESMLE(S1)	CESMLE(S2)	CESMLE(S3)
N	10	10	11.09	7.03	4.24	2.26
Mean	8	-1.1	7.80	8.45	6.77	5.34
SD	47	19	43.18	43.66	35.38	33.88
10 th P	-44.95	-6.01	-45.00	-49.07	-38.74	-22.82
Q1	-5.04	0.00	-3.51	0.00	0.00	0.00
Median	0	0	0.00	0.00	0.00	0.00
Q2	18.16	0.00	28.02	32.72	6.21	10.28
90 th P	82.38	3.58	84.30	81.65	63.00	45.00
¹ Normality p	0.18	<0.01*	<0.01*	<0.01*	<0.01*	<0.01*
² Welch p		0.58	0.99 0.19	0.86	0.78	0.58
³ Tukey p		1.00	1.00 0.98	1.00	1.00	1.00
⁴ WMW p		0.90	0.68 0.52	0.41	1.00	0.83

All tests significant at $\alpha = 0.05$.

¹ Shapiro-Wilk Normality Test. * denotes the distribution is not a normal distribution.

² Welch Two-Sample T-Test. * denotes the paired sample distribution means are significantly different.

³ Tukey Honestly Significant Difference Test. * denotes the paired sample distribution means are significantly different.

⁴ Wilcoxon Mann-Whitney U Test. * denotes the sample distributions are significantly different.