

Table: Statistics for variable (axis-ang) and region (wna-s). Model N is the average 34-member objects.

Statistic	CRU	PRISM	CESMLE(C)	CESMLE(S1)	CESMLE(S2)	CESMLE(S3)
N	17	10	7.03	7.15	8.38	7.03
Mean	14	-5.7	2.47	13.18	-0.43	-1.70
SD	42	50	55.17	55.06	51.18	42.62
10 th P	-44.54	-63.88	-69.43	-62.45	-68.70	-53.96
Q1	0.00	-40.36	-45.74	-30.12	-38.64	-24.82
Median	0	0	0.00	0.00	0.00	-0.60
Q2	50.86	0.00	46.30	66.84	33.29	15.63
90 th P	65.88	51.45	90.00	90.00	90.00	57.91
¹ Normality p	0.06	0.26	< 0.01 *	< 0.01 *	< 0.01 *	< 0.01 *
² Welch p		0.31	0.29 0.63	0.03 *	0.54	0.36
³ Tukey p		0.93	0.94 1.00	0.20	0.99	0.95
⁴ WMW p		0.27	0.37 0.70	0.04 *	0.55	0.29

All tests significant at $\alpha = 0.05$.

¹ Shapiro-Wilk Normality Test. * denotes the distribution is not a normal distribution.

² Welch Two-Sample T-Test. * denotes the paired sample distribution means are significantly different.

³ Tukey Honestly Significant Difference Test. * denotes the paired sample distribution means are significantly different.

⁴ Wilcoxon Mann-Whitney U Test. * denotes the sample distributions are significantly different.