

Table: Statistics for variable (axis-ang) and region (wna-s). Model N is the average 34-member objects.

Statistic	CRU	PRISM	CESMLE(C)	CESMLE(S1)	CESMLE(S2)	CESMLE(S3)
N	21	13	7.94	5.44	3.12	2.29
Mean	19	13	13.39	12.71	13.41	23.43
SD	42	46	55.38	52.68	51.36	56.40
10 th P	-21.15	-51.90	-62.82	-58.60	-55.39	-59.68
Q1	0.00	0.00	-25.47	-14.08	-9.15	0.00
Median	0	15	0.00	0.00	0.00	0.00
Q2	47.76	49.22	69.40	59.10	59.95	90.00
90 th P	90.00	63.50	90.00	90.00	90.00	90.00
¹ Normality p	0.01*	0.66	<0.01*	<0.01*	<0.01*	<0.01*
² Welch p		0.73	0.58 1.00	0.90	1.00	0.17
³ Tukey p		1.00	1.00 1.00	1.00	1.00	0.69
⁴ WMW p		0.90	0.60 0.87	0.91	0.94	0.07

All tests significant at $\alpha = 0.05$.

¹ Shapiro-Wilk Normality Test. * denotes the distribution is not a normal distribution.

² Welch Two-Sample T-Test. * denotes the paired sample distribution means are significantly different.

³ Tukey Honestly Significant Difference Test. * denotes the paired sample distribution means are significantly different.

⁴ Wilcoxon Mann-Whitney U Test. * denotes the sample distributions are significantly different.